



New P-T-t data on the metamorphic sole of the Amasia ophiolites and implications for the geodynamical process, NW of the Sevan-Akera suture zone, Lesser Caucasus (Armenia)

Hässig Marc, Yann Rolland, Marc Sosson, Ghazar Galoyan

► To cite this version:

Hässig Marc, Yann Rolland, Marc Sosson, Ghazar Galoyan. New P-T-t data on the metamorphic sole of the Amasia ophiolites and implications for the geodynamical process, NW of the Sevan-Akera suture zone, Lesser Caucasus (Armenia). EGU General Assembly 2011, Apr 2011, Vienne, Austria. pp. EGU2011-7462-1. hal-01060852

HAL Id: hal-01060852

<https://hal.science/hal-01060852>

Submitted on 12 Sep 2014

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



New P-T-t data on the metamorphic sole of the Amasia ophiolites and implications for the geodynamical process, NW of the Sevan-Akera suture zone, Lesser Caucasus (Armenia)

Marc Hässig (1), Yann Rolland (1), Marc Sosson (1), and Ghazar Galoyan (2)

(1) Université de Nice-Sophia Antipolis, Lab. GEOAZUR - OCA, France (hassig@geoazur.unice.fr), (2) Institute of Geological Sciences, National Academy of Sciences of Armenia, Armenia

The ophiolites of the Lesser Caucasus belong to the Tethyan Ophiolitic Belt. In the northwestern part of the Sevan Akera suture zone (Lesser Caucasus, Armenia) ophiolites correspond to a major obduction of oceanic crust over the South Armenian continental block. Near the locality of Amasia (NW Armenia), our mapping evidenced a series of (1) unmetamorphosed gabbroic oceanic crust, resting on (2) serpentinites and a greenschist grade tectonic mélange composed of deformed pillow-basalts, radiolarites and cherts, and (3) a basal slice of garnet amphibolites bearing similar compositional features as the ophiolite.

They are sliced by post Eocene thrusts related to the shortening of the suture after the collision of the South Armenian Block (Sosson et al., 2010). Structural observations allow us to consider these metamorphic rocks (garnet amphibolites and quartz-garnet-oligoclase-amphibole-chlorite schistes) to be the sole of the obducted ophiolite.

The metamorphic rocks allow the evaluation of pressure-temperature-time conditions endured by the base of the oceanic crust during the obduction of the Armenian ophiolite series. Pressure and temperature estimations were obtained by calculating equilibrium conditions between different mineral phases which coexist in the sample coupled with whole rock geochemistry using the software PERPLEX. The P-T data show a counterclockwise path with (1) a HT-LP peak of $P = 6-7$ kbar and $T > 630^{\circ}\text{C}$, followed by MP-MT peak at $P = 8-10$ kbar and $T = 600^{\circ}\text{C}$. In order to further constrain the evolution of the amphibolites and build a P-T-t path using $40\text{Ar}/39\text{Ar}$ dating on amphiboles and white micas (phengite). The ages obtained on both HT and MT peaks are similar within error ($90.8 \text{ Ma} \pm 3.0$ (2σ), and $90.8 \text{ Ma} \pm 1.2$ (2σ), respectively), which suggests a very rapid cooling of the metamorphic sole during its stacking in the subduction zone.

The counterclockwise path determined confirms the observations made below the Oman ophiolite by Goffé et al. (1988). This path is also in agreement with a scaling process of oceanic crust during the obduction. In this scenario slices of oceanic crust would have been dragged under the obduction front below a relatively hot oceanic crust. Afterwards, these slices would then have been amphibolitized along a HT-MP geothermal gradient. Rapidly, a thickening of the system at c. 91-90 Ma, and underthrusting of amphibolites would have caused the conditions to change to MP-MT. This infers that the South Armenian Block enters the subduction zone at this date.

A recent study has shown metamorphic rocks in the same structural setting were affected by a HP metamorphism in the Blue Schist facies (Rolland et al., 2009). The age of this blue schist of Stepanavan, Armenia is slightly older (95-91 Ma) (Galoyan et al., 2009) than the amphibolites of Amasia (91-90 Ma). This assumes the occurrence of an active subduction just before the development of the HT-MP metamorphism of Amasia.

References:

Galoyan, G., Rolland, Y., Sosson, M., Corsini, M., Billo, S., Verati, C., Melkonyan, R., 2009. Geology, geochemistry and $40\text{Ar}/39\text{Ar}$ dating of the Sevan ophiolites (Lesser Caucasus, Armenia): Evidence for Jurassic Back-arc opening and hot spot event between the South Armenian Block and Eurasia. *Journal of Asian Earth Science*, 34, 135-153.

Goffé, B., Michard, A., Kienast, J.R., Le Mer, O., 1988. A case of obduction-related high-pressure, low-temperature metamorphism in upper crustal nappes, Arabian continental margin, Oman: P-T paths and kinematic interpretation. *Tectonophysics*, 151, 363-386.

Rolland, Y., Billo, S., Corsini, M., Sosson, M., Galoyan, G., 2009. Blueschists of the Amassia-Stepanavan Suture Zone (Armenia): linking Tethys subduction history from E-Turkey to W-Iran. *International Journal of Earth Science*, 98, 533-550.

Sosson, M., Rolland, Y., Müller, C., Danelian, T., Melkonyan, R., Kekelia, S., Adamia, S., Badazadeh, V., Kangarli, T., Avagyan, A., Galoyan, G., Mosar, J., 2010. Subduction, obduction and collision in the Lesser Caucasus (Armenia, Azernaijan, Georgia), new insights. *Geological Society, London, Special Publications*, 340, 329-352.